

FIG. 1

008027" 95T2260





FIG. 2

FIG.3



Plaque reduction assay with serum from CD4 T-cell
knockout mice immunized with inactivated PR8
viruses




FIG. 4




FIG. 5A




FIG. 5B




FIG. 5C

The figure consists of three vertically stacked bar charts, each representing a different antibody isotype: IgM (top), IgG2a (middle), and IgG3 (bottom). The y-axis for all charts is 'Concentration (ug/ml)'.

- IgM Chart:** The y-axis ranges from 0 to 50. The 'Con.' group has bars for 'First' (~4.5) and 'Boost' (~4.0). The 'PR8' group has bars for 'First' (~14.0) and 'Boost' (~16.5). Error bars are present on all bars.
- IgG2a Chart:** The y-axis ranges from 0 to 20. The 'Con.' group has bars for 'First' and 'Boost' that are very close to zero. The 'PR8' group has bars for 'First' and 'Boost' that are also very close to zero.
- IgG3 Chart:** The y-axis ranges from 0 to 20. The 'Con.' group has bars for 'First' and 'Boost' that are very close to zero. The 'PR8' group has bars for 'First' and 'Boost' that are very close to zero.

Legend for all charts: □ First, ■ Boost.




FIG. 6

U.S. DEPARTMENT OF THE INTERIOR




FIG. 7

00802T " 99T E 60 CD8




FIG. 8